

Облікова картка дисертації

I. Загальні відомості

Державний обліковий номер: 0414U003945

Особливі позначки: відкрита

Дата реєстрації: 14-10-2014

Статус: Захищена

Реквізити наказу МОН / наказу закладу:



II. Відомості про здобувача

Власне Прізвище Ім'я По-батькові:

1. Новохацька Ольга Василівна

2. Olga Novokhatska

Кваліфікація:

Ідентифікатор ORCID ID: Не застосовується

Вид дисертації: кандидат наук

Аспірантура/Докторантура: так

Шифр наукової спеціальності: 03.00.03

Назва наукової спеціальності: Молекулярна біологія

Галузь / галузі знань: Не застосовується

Освітньо-наукова програма зі спеціальності: Не застосовується

Дата захисту: 30-09-2014

Спеціальність за освітою: 7.070408

Місце роботи здобувача: Інститут молекулярної біології і генетики НАН України

Код за ЄДРПОУ: 05417101

Місцезнаходження: 03680 Київ, вул. Заболотного, 150

Форма власності:

Сфера управління: Національна академія наук України

Ідентифікатор ROR: Не застосовується

III. Відомості про організацію, де відбувся захист

Шифр спеціалізованої вченої ради (разової спеціалізованої вченої ради): Д26.237.01

Повне найменування юридичної особи:

Код за ЄДРПОУ:

Місцезнаходження:

Форма власності:

Сфера управління:

Ідентифікатор ROR: Не застосовується

IV. Відомості про підприємство, установу, організацію, в якій було виконано дисертацію

Повне найменування юридичної особи: Інститут молекулярної біології і генетики НАН України

Код за ЄДРПОУ: 05417101

Місцезнаходження: 03680 Київ, вул. Заболотного, 150

Форма власності:

Сфера управління: Національна академія наук України

Ідентифікатор ROR: Не застосовується

V. Відомості про дисертацію

Мова дисертації:

Коди тематичних рубрик: 34.15.25

Тема дисертації:

1. Характеристика адаптерного білка інтерсектину 2 людини
2. Characterization of human adaptor protein intersectin 2

Реферат:

1. Інтерсектин 1 та 2 людини - ендцитозні адаптерні білки, що характеризуються високим ступенем гомології, проте асоційовані із різними захворюваннями людини. Для того щоб визначити, наскільки інтерсектин 1 та 2 можуть бути функціонально розділеними, було проведено порівняльний аналіз цих білків. З огляду на диференційні рівні експресії ITSN2 у ембріональних тканинах людини нами було досліджено участь ITSN2 у ранньому ембріональному розвитку. Встановлено, що надекспресія С-кінцевого регіону білка, специфічного для довгої ізоформи, викликає дефекти розвитку ембріонів *Xenopus laevis*, формування яких пов'язане з порушеннями структури актинового цитоскелету. Показано, що інтерсектин 1 та 2 колокалізуються в клітині та формують комплекс. Порівняння структури доменів SH3 інтерсектинів показало подібність їх поверхонь зв'язування. Встановлено, що домени SH3 ITSN2 можуть взаємодіяти із білками-партнерами ITSN1, а також ідентифікованими новими білками-партнерами інтерсектинів. Виявлено новий інтерфейс взаємодії, представлений тирозиновими мотивами в ITSN2. В різних лініях клітин людини було показано фосфорильовання ізоформ ITSN2. Стимуляція епідермальним фактором росту призводить до підвищення

рівня фосфорилування ITSN2 та його впізнавання SH2-вмісними білками, а саме кіназами Fyn, Fgr та Abl1, регуляторною субодиницею PI3K, адаптерними білками Grb2 та Crk, а також фосфоліпазою C?. Отже, під час еволюції хребетних ITSN2 набув нового інтерфейсу для білок-білкових взаємодій, що забезпечило його специфічне впізнавання доменами SH2 сигнальних білків. Ці дані можуть бути важливими для розуміння функціонального різноманіття білків-паралогів родини інтерсектинів та їх ролі у захворюваннях людини. Ключові слова: інтерсектин, ранній ембріональний розвиток, білок-білкові взаємодії, домени SH3, фосфорилування за залишками тирозину.

2. Modern complexity of the eukaryotic membrane transport system is considered the result of paralogous gene expansion. In mammals clathrin-mediated endocytosis (CME) is mediated by a machinery of proteins represented mostly by at least two paralogues. Scaffolding proteins of the intersectin (ITSN) family, ITSN1 and ITSN2, are crucial for the initiation stage of clathrin-mediated endocytosis. These proteins are closely related but have implications in distinct pathologies. So far it remains obscure to what extent the roles of ITSNs are common in tissues where they are coexpressed. Thus, the aim of this work was to characterize ITSN2 and perform a comparative analysis with ITSN1. Given differential expression of ITSN2 long isoform in human embryonic tissues we studied its involvement in early embryonic development using *Xenopus laevis* as a model. We show that transcripts of ITSN2-L can be detected in *Xenopus* embryos from the first cleavage onwards. Overexpression of functional domains of ITSN2-L in embryos resulted in aberrant phenotypes. The strongest phenotype was produced by the C-terminal extension of ITSN2-L. Embryos displayed hyperpigmentation and gastrulation failure that were incompatible with survival. The C-terminus of ITSN2-L includes the DH-PH tandem, a nucleotide exchange factor for the small GTPase Cdc42 and the C2 domain. Further investigations revealed that the DH-PH tandem was responsible for the development of the phenotype affecting the actin cytoskeleton in embryos. Observed developmental defects depended on Cdc42. The effect of expression of the constitutively active GTPase strongly resembled that of the DH-PH tandem. The dominant negative Cdc42 partially rescued developmental defects induced by the expression of the DH-PH tandem. Thus, our data indicate that the ITSN2 exchange factor regulates the activity of Cdc42 during embryo development affecting actin cytoskeleton in *Xenopus* embryos. The next step of this work was functional comparison of intersectins. So far these proteins were investigated independently except for a recent study of Henne et al. However, a comparison of ITSN1 and ITSN2 is of current importance to understand to what extent they could substitute for each other under normal and pathological conditions. The data about ITSN2 and its functional role are limited. It was shown that many paralogous proteins involved in membrane-trafficking processes perform similar functions in distinct cellular compartments. We now provide evidence that ITSN1 and its paralogue ITSN2 are components of common protein complexes and have a predominantly common intracellular distribution in the cell line studied. Previously, ITSN1 was suggested to be constitutively associated with the endocytic adaptor Eps15 in a protein complex. Presumably, ITSN2 is also engaged in this scaffolding complex crucial for progression of clathrin-mediated endocytosis. Regarding the fact that ITSNs have the same domain organization, a similar subcellular distribution and are components of common complexes, we investigated to what extent pools of their binding partners could be different. Most known ITSN1 interactors are ligands of its five SH3 domains. A comparison of the SH3 domain structures of ITSNs showed a high similarity of their ligand-binding sites. Amino acid mismatches within respective SH3 domain pairs were predominantly located beyond the ligand-binding sites. The results from in vitro binding experiments demonstrate the ability of the SH3 domains of ITSN2 to pull down all the protein partners of ITSN1 investigated here. Moreover, novel SH3 ligands, the endocytic adaptors POB1 and Repl1 as well as the signaling proteins Sema6A and SPRY2, were common to both ITSNs. Data from the comparison of ligand-binding sites and in vitro binding experiments suggest that the SH3 domains of ITSNs bind predominantly similar ligands. Certainly, the possibility of specific partners for the certain SH3 domains can not be excluded. There are several possible explanations for this phenomenon. First, selection kept the ligand binding sites of the ITSNs SH3 domains similar whereas amino acid residues beyond the sites varied to a greater extent. Another explanation concerns noncanonical functions of the SH3 domains and the existence of additional interaction interfaces. Given identified similarities between the ITSNs, we searched for novel binding interfaces that could provide specific protein-protein interactions. Sequence

analysis revealed that ITSNs differ in the number of tyrosine residues and their location within the molecule. We intended to search for a putative novel protein–interaction interface that could be distinct between ITSN molecules. A comparison of the primary sequences of the ITSNs demonstrated that ITSN2-S contains 60% more tyrosine residues than ITSN1-S. Accumulation of tyrosines in ITSN2 isoforms was clearly observed in the range of its vertebrate orthologues from fish to primates. All the conserved tyrosine residues of ITSN1-S were located within protein domains whereas additional tyrosine residues in ITSN2-S were found in interdomain regions. Tyrosine phosphorylation of ITSN2 isoforms but not ITSN1-S was detected in various cell lines. This is in good correspondence with phosphoproteomic data about this modification of ITSNs. Acquisition by ITSN2 of additional tyrosine residues and their posttranslational modification lead us to assume that tyrosinebased linear motifs emerged during evolution to regulate ITSN2. Evolution of linear motifs is considered to be the fastest and major mechanism involved in changing protein interaction networks. One could expect that phosphorylation of the linear motif allows recognition of the motif by proteins bearing phosphotyrosine-binding domains. We have demonstrated that the SH2 domains of the kinases Fyn, Fgr and Abl1, the regulatory subunit of PI3K, the adaptor proteins Grb2 and Crk, and phospholipase C gamma could mediate binding to ITSN2. In spite of phosphorylation of ITSN2 isoforms in growing cells, interactions with the SH2 domains were detected only in EGF-stimulated cells. It could be suggested that EGF treatment induces specific phosphorylation of motifs recognized by certain SH2 domains. Our results indicate that during evolution of vertebrates ITSN2 acquired novel protein–interaction interface that allows its specific recognition by the SH2 domains of signaling proteins. We propose that these data could be important for understanding the functional diversity of paralogous ITSN proteins and establishing their roles in diseases. Keywords: intersectin, early embryonic development, protein–protein interactions, SH3 domains, tyrosine phosphorylation.

Державний реєстраційний номер ДіР:

Пріоритетний напрям розвитку науки і техніки:

Стратегічний пріоритетний напрям інноваційної діяльності:

Підсумки дослідження:

Публікації:

Наукова (науково–технічна) продукція:

Соціально–економічна спрямованість:

Охоронні документи на ОПВ:

Впровадження результатів дисертації:

Зв'язок з науковими темами:

VI. Відомості про наукового керівника/керівників (консультанта)

Власне Прізвище Ім'я По–батькові:

1. Риндич Алла Володимирівна
2. Alla Rynditch

Кваліфікація: д.б.н., 03.00.03

Ідентифікатор ORCID ID: Не застосовується

Додаткова інформація:

Повне найменування юридичної особи:

Код за ЄДРПОУ:

Місцезнаходження:

Форма власності:

Сфера управління:

Ідентифікатор ROR: Не застосовується

VII. Відомості про офіційних опонентів та рецензентів

Офіційні опоненти

Власне Прізвище Ім'я По-батькові:

1. Скок Марина Володимирівна

2. Скок Марина Володимирівна

Кваліфікація: д.б.н., 03.00.03

Ідентифікатор ORCID ID: Не застосовується

Додаткова інформація:

Повне найменування юридичної особи:

Код за ЄДРПОУ:

Місцезнаходження:

Форма власності:

Сфера управління:

Ідентифікатор ROR: Не застосовується

Власне Прізвище Ім'я По-батькові:

1. Сиволоб Андрій Володимирович

2. Сиволоб Андрій Володимирович

Кваліфікація: д.б.н., 03.00.03

Ідентифікатор ORCID ID: Не застосовується

Додаткова інформація:

Повне найменування юридичної особи:

Код за ЄДРПОУ:

Місцезнаходження:

Форма власності:

Сфера управління:

Ідентифікатор ROR: Не застосовується

Рецензенти

VIII. **Заключні відомості**

**Власне Прізвище Ім'я По-батькові
голови ради**

Єльська Ганна Валентинівна

**Власне Прізвище Ім'я По-батькові
головуючого на засіданні**

Єльська Ганна Валентинівна

**Відповідальний за підготовку
облікових документів**

Реєстратор

**Керівник відділу УкрІНТЕІ, що є
відповідальним за реєстрацію наукової
діяльності**



Юрченко Т.А.